




WEB of SCIENCE Core Collection

Universitatea de Medicină și Farmacie „Grigore T. Popa”, Iași *
2010-2020


WEB of SCIENCE Core Collection, UMF IS, 2010-2020



Highly Cited Papers	24
Review Articles	597
Early Access	1
Open Access	1,530
Associated Data	20


Highly Cited Paper

A paper that belongs to the top 1% of papers in a research field published in a specified year.




WEB of SCIENCE Core Collection, UMF IS, 2010-2020

Distribution on publication years based




WEB of SCIENCE Core Collection, UMF IS, 2010-2020

Distribution on Document Types based



WEB of SCIENCE Core Collection, UMF IS, 2010-2020

Distribution on Web of Science Index based




WEB of SCIENCE Core Collection, UMF IS, 2010-2020

Web of Science Categories (top 25)



WEB of SCIENCE Core Collection, UMF IS, 2010-2020

Research Areas (top 25)




WEB of SCIENCE Core Collection, UMF IS, 2010-2020

Sources (journals, proceedings) (Top 25)




WEB of SCIENCE Core Collection, UMF IS, 2010-2020

Countries co-authorship (Top 24)





WEB of SCIENCE Core Collection, UMF IS, 2010-2020

Institutions co-authorship (Top 24)





WEB of SCIENCE Core Collection, UMF IS, 2010-2020


Most cited papers (Top 5)

- | | | |
|----------------------------|--|---|
| <input type="checkbox"/> 1 | 

Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults
Ezzati, M; Bentham, J; (...); Cisneros, JZ
Dec 16 2017 LANCET 390 (10113) , pp.2627-2642



Background Underweight, overweight, and obesity in childhood and adolescence are associated with adverse health consequences throughout the life-course. Our aim was to estimate worldwide trends in mean body-mass index (BMI) and a comprehensive set of BMI categories that cover underweight to obesity in children and adolescents, z ... Show more

Free Full Text From Publisher *** | 2,407
Citations
<hr/> 54
References
<hr/> Related records |
| <input type="checkbox"/> 2 | 

Towards the introduction of the 'Immunoscore' in the classification of malignant tumours
Galon, J; Mlecnik, B; (...); Pages, F
Jan 2014 JOURNAL OF PATHOLOGY 232 (2) , pp.199-209



The American Joint Committee on Cancer/Union Internationale Contre le Cancer (AJCC/UICC) TNM staging system provides the most reliable guidelines for the routine prognostication and treatment of colorectal carcinoma. This traditional tumour staging summarizes data on tumour burden (T), the presence of cancer cells in draining and regional lymph n ... Show more

Free Full Text From Publisher *** | 803
Citations
<hr/> 70
References
<hr/> Related records |
| <input type="checkbox"/> 3 | 
International validation of the consensus Immunoscore for the classification of colon cancer: a prognostic and accuracy study
Pages, F; Mlecnik, B; (...); Galon, J
May 26 2018 LANCET 391 (10135) , pp.2128-2139

Background The estimation of risk of recurrence for patients with colon carcinoma must be improved. A robust immune score quantification is needed to introduce immune parameters into cancer classification. The aim of the study was to assess the prognostic value of total tumour-infiltrating T-cell counts and cytotoxic tumour-infiltrating T-cells cou ... Show more

Full Text at Publisher *** | 728
Citations
<hr/> 33
References
<hr/> Related records |
| <input type="checkbox"/> 4 | 

Fibrinolysis for Patients with Intermediate-Risk Pulmonary Embolism
Meyer, G; Vicaut, E; (...); Konstantinides, SV
Apr 10 2014 NEW ENGLAND JOURNAL OF MEDICINE 370 (15) , pp.1402-1411

BackgroundThe role of fibrinolytic therapy in patients with intermediate-risk pulmonary embolism is controversial. MethodsIn a randomized, double-blind trial, we compared tenecteplase plus heparin with placebo plus heparin in normotensive patients with intermediate-risk pulmonary embolism. Eligible patients had right ventri ... Show more

Free Published Article From Repository Full Text at Publisher *** | 701
Citations
<hr/> 28
References
<hr/> Related records |
| <input type="checkbox"/> 5 | 

EuReCa ONE-27 Nations, ONE Europe, ONE Registry A prospective one month analysis of out-of-hospital cardiac arrest outcomes in 27 countries in Europe
Grasner, JT; Lefering, R; (...); Bossaert, LL
Aug 2016 RESUSCITATION 105 , pp.188-195

Introduction: The aim of the EuReCa ONE study was to determine the incidence, process, and outcome for out of hospital cardiac arrest (OHCA) throughout Europe.
Methods: This was an international, prospective, multi-centre one-month study. Patients who suffere ... Show more | 390
Citations
<hr/> 29
References |